

PRODUCT INFORMATION

Tag	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 C-Flag&Strep Tag
Target	TRPM5
Synonyms	LTRPCS, MTR1
Description	Human TRPM5-Strep full length protein-synthetic nanodisc
Delivery	6-8weeks
Uniprot ID	Q9NZQ8
Expression Host	HEK293
Protein Families	Ion Channels: Transient receptor potential
Protein Pathways	N/A
Molecular Weight	The human full length TRPM5-Strep protein has a MW of 131.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the transient receptor potential (TRP) protein family, which is a diverse group of proteins with structural features typical of ion channels. This protein plays an important role in taste transduction, and has characteristics of a calcium-activated, non-selective cation channel that carries Na ⁺ , K ⁺ , and Cs ions equally well, but not Ca ²⁺ ions. It is activated by lower concentrations of intracellular Ca ²⁺ , and inhibited by higher concentrations. It is also a highly temperature-sensitive, heat activated channel showing a steep increase of inward currents at temperatures between 15 and 35 degrees Celsius. This gene is located within the Beckwith-Wiedemann syndrome critical region-1 on chromosome 11p15.5, and has been shown to be imprinted, with exclusive expression from the paternal allele. [provided by RefSeq, Oct 2010]
Usage	Research use only
Conjugate	Unconjugated

