

**PRODUCT INFORMATION**

<b>Target</b>	CXADR
<b>Synonyms</b>	CAR;MCAR;MCVADR
<b>Description</b>	Recombinant mouse CXADR protein with C-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P97792
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-6×His Tag
<b>Molecular Characterization</b>	Mouse CXADR(Leu20-Gly237) 6×His tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 24.9 kDa after removal of the signal peptide. The apparent molecular mass of mCXADR-His is approximately 15-35 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Yefei_Storage</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a protein that is part of the Cortical Thymocyte marker in Xenopus (CTX) subfamily within the immunoglobulin superfamily. Members of this subfamily, predominantly expressed on the surface of endothelial and epithelial cells, help establish cell polarity and provide a barrier function, regulating migration of immune cells. This protein, first identified as the receptor for adenovirus subgroup C and coxsakieviruses group B, is developmentally regulated and plays an important role in cardiac development. Alternative splicing results in multiple transcript variants that encode different protein isoforms. [provided by RefSeq, Jan 2013]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated





Figure 1. Mouse CXADR Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

