Cat. No. DMC101357P



PRODUCT INFORMATION

Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products-public.php on line 2806 1506 Clone ID

Synonyme Host Species

Description

Delivery

Uniprot ID

IgG type

Clonality Reactivity Applications Recommended Dilutions

Purification

Formulation & Reconstitution

Storage & Shipping

This gene encodes a receptor that mediates the functions of activins, which are members of the transforming growth factor-beta (TGF involved in diverse biological processes. The encoded protein is a transmembrane serine-threonine kinase receptor which mediates is heterodimeric complexes with various combinations of type I and type II receptors and ligands in a cell-specific manner. The encoded primarily involved in ligand-binding and includes an extracellular ligand-binding domain, a transmembrane domain and a cytoplasmic kinase domain. This gene may be associated with susceptibility to preeclampsia, a pregnancy-related disease which can result in materioribidity and mortality. Alternative splicing results in multiple transcript variants of this gene. [provided by RefSeq. Jun 2013] Research use only Background

Conjugate

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

