

## **PRODUCT INFORMATION**

Clone ID DM180 CCR8 **Target** 

CC-CKR-8; CCR-8; CDw198; CKRL1; CMKBR8; CMKBRL2; CY6; GPRCY6; TER1 **Synonyms** 

**Host Species** 

PE-conjugated Anti-CCR8 antibody(DM180); **Description** 

Rabbit mAb Delivery 3-4 weeks **Uniprot ID** P51685 IgG type Rabbit IgG

Clonality Monoclonal Reactivity Human

Recommended

**Applications** 

Formulation &

Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Liquid
☐PBS with 0.05% Proclin300, 1% BSA Reconstitution

Flow Cyt

Storage & Shipping Store at 2°C-8°C for 6 months

> This gene encodes a member of the beta chemokine receptor family; which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptors are important for the migration of various cell types into the inflammatory sites. This receptor protein preferentially expresses in the thymus. I-309; thymus activation-regulated cytokine (TARC) and macrophage inflammatory protein-1 beta (MIP-1 beta) have been identified

**Background** as ligands of this receptor. Studies of this receptor and its ligands suggested its role in regulation of monocyte chemotaxis and thymic cell apoptosis. More specifically; this receptor may contribute to the proper positioning of activated T cells within the antigenic challenge sites and specialized areas of lymphoid tissues.

This gene is located at the chemokine receptor gene cluster region.

Usage Research use only Conjugate PE-conjugated


All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under

patent application. Any protein sequencing or **DIMA Disclaimer** reverse engineering attempt is prohibited. We are

actively scrutinizing all patent application to

Email: info@dimabio.com Website: www.dimabio.com

ensure no IP infringement.



/+86-400-006-0995(China)

