

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 bM165 Clone ID

CLEC12A

CLEC12A;MICL;CLL-1;CLL1;DCAL2;DCAL-2;CD371 Synonyme

Host Species Rabbit

PE-conjugated Anti-CLEC12A antibody(DM165); Rabbit mAb Description

Delivery Under Development Uniprot ID Q5QGZ9 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human Flow Cyt Application

Recommended Dilutions Flow Cyt 1:100 Purification

Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Liquid PBS with 0.05% Proclin300, 1% BSA

Storage & Shipping Store at 2°C-8°C for 6 months

This gene encodes a member of the C-type lectin-C-type lectin-like domain (CTL:CTLD) superfamily. Members of this family share a common protein fold and have diverse functions; such as cell adhesion; cell-cell signaling, glycoprotein turnover; and roles in inflammation and immune response. The protein encoded by this gene is a negative regulator of granulocyte and monocyte function. Several alternatively spliced transcript variants of this gene have been described; but the full-length nature of some of these variants has not been determined. This gene is closely linked to other CTL:CTLD superfamily members in the natural killer gene complex region on chromosome 12p13. Background

Usage Research use only Conjugate

Research use unity
PE-conjugated
All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. Actin sequ. DIMA Disclaimer

Email: info@dimabio.com Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)